



Bicornuate Uterus With Multiple Uterine Fibroids, Metroplasty and Myomectomy; Case Profile and Literature Review

Akani CI¹; John CO¹; Korubo I¹; Omoruyi S; Eli S²; Akani UE³, Olaka EW¹

Department of Obstetrics and Gynaecology, University of Port Harcourt Teaching Hospital.¹
Mother, Baby and Adolescent Care Global Foundation.²

Department of Public Health, University of Port Harcourt Teaching Hospital.³

ARTICLE INFO

Article No.:022622023

Type: Case Study

Full Text: [PDF](#), [HTML](#), [EPUB](#), [PHP](#)

Accepted: 28/02/2022

Published: 25/03/2022

***Corresponding Author**

Prof C.I Akani MBBS, FWACS, FICS

E-mail: ciakaniph@gmail.com

Phone: 08103011111

Keywords: Bicornuate uterus, uterine fibroid, metroplasty, myomectomy.

INTRODUCTION

The prevalence of uterine malformations like bicornuate, septate or arcuate uterus in the general population is about 6.7% but in patients with recurrent miscarriage it is about 16.7%.¹ Bicornuate uterus is one of the most commonly diagnosed mullerian duct anomaly, constituting 25% of all uterine anomalies and it is a type

IV mullerian duct anomaly according to the American Society of Reproductive Medicine.^{2,3} Other classes include I-Hypoplasia/Agenesis, II-Unicornuate, III-Didelphus, V-septate, VI-arcuate and VII-Diethylstilbestrol drug related.^{2,3}




Figure 1

EMBRYOLOGY AND PATHOPHYSIOLOGY

The class IV malformation (Bicornuate uterus) is caused by partial non-fusion of the upper part of the müllerian ducts. This results in a central myometrium that may extend to the level of the internal cervical os (bicornuate unicollis) or external os (bicornuate bicollis), with a fundal cleft 1cm deep.³⁻⁶ The horns of the bicornuate uteri are not as fully developed and are smaller than those in the didelphus uteri.

CLINICAL FEATURES

Patients with bicornuate uterus are usually asymptomatic but can present with symptoms like menorrhagia and dysmenorrhea which are non-specific symptoms and also a history of recurrent miscarriage, preterm deliveries and persistent abnormal lies and presentation in pregnancy.²⁻⁴ The diagnosis is usually made as an incidental finding during evaluation for infertility and patients with recurrent miscarriage.^{2-4,7}

INVESTIGATIONS

It is important to differentiate a bicornuate from a septate uterus. Hysterosalpingogram (HSG) alone cannot differentiate these entities, because this imaging approach cannot evaluate the external contour of the uterus.⁸ While laparoscopy was used primarily for this purpose in the past, modern imaging techniques including 3D ultrasonography and MRI can adequately differentiate these two entities. Imaging criteria to differentiate septate and bicornuate uteri have been developed. A septate uterus has a flat or convex fundus

or a fundal indentation 60°.^{3,8} On MRI, a septate uterus will fail to show an intervening myometrium between the T2-hypointense septum that separates the endometrial cavities [80, 81]. In contrast, a bicornuate uterus will show two T2-hyperintense endometrial cavities, each with a junctional zone and myometrial band of intermediate signal intensity.

Endoscopic procedures like laparoscopy and hysteroscopy are diagnostic and therapeutic.

MANAGEMENT

Most cases of Bicornuate uterus may not need any treatment unless they are associated with infertility, recurrent pregnancy loss or Uterine pathologies like fibroids.^{6,8}

Conventional transabdominal metroplasty has been shown to significantly improve the pregnancy outcome in patients with bicornuate uterus.^{2,8} Laparoscopic approach is also technically challenging but offers the general positive benefits of endoscopic surgeries. Thus, the most common surgical treatment options for bicornuate uterus may include the Strassman metroplasty and cervical cerclage. The surgery entails removing the abnormal tissue that separates the cornua of the uterus, then using several layers of stitches to create a normal shape and single uterine cavity. The pregnancy rate following metroplasty has been seen in up to 90% of cases.⁴

CASE REPORT

A 35 year old nulliparous woman who presented to the gynaecological clinic with complaints of recurrent lower abdominal pains, dysmenorrhea and abdominal mass of 3 years duration. There was no menorrhagia, urinary or pressure symptoms from the mass.

Abdominal examination revealed a 22 week sized abdomeninopelvic mass. She was evaluated for

symptomatic uterine fibroids. Incidentally the hysterosalpingography revealed a congenitally malformed uterus, suggested to be bicornuate uterus. She was further assessed using a diagnostic hysteroscopy which showed the obvious septation with the two cavities. An intravenous urography ruled out a pathology of the urinary system. These are shown in figure 2.





Figure 2: (A)-HSG showing bicornuate uterus. (B)findings of septal protrusion on hysteroscopy

A diagnosis of symptomatic uterine fibroids co-existing with a bicornuate uterus was made. She was counseled and taken up for Abdominal Myomectomy and Metroplasty for uterine fibroids in a Bicornuate uterus.

Intra-operatively, the abdomen was opened by a midline incision. The uterus was exteriorized and inspected to confirm the two horns with obvious big fibroid nodules. The tubes and ovaries were normal. A conventional myomectomy was done using only anterior uterine wall incision. An incision extended from the


superior aspect of each horn near the interstitial region of the fallopian tubes to the inferior aspect of the uterus was made to access the two cavities. The endometrium of both cavities were exposed, septum was identified and excised. Apposition of the myometrium excluding endometrium was done using interrupted sutures with 2-0 PDS to form a single uterine cavity. The rest of the uterus was reconstituted using conventional surgical techniques. The uterus was reperitonized and the abdomen was closed. The next figure below; figure 3, shows intra-operative steps.




A-Uterus with fibroids on the two horns




B- The uterine horns and appendages shown (grasped with the two babcock forceps)



C- Yellow and Green tags showing the two different cavities, blue line showing the septum.



Uterine septum coloured yellow above



D- Septum excised and a single cavity with catheter bulb in situ



E- Reconstructed uterus

Figure 3: Intra-operative steps (A,B,C,D, & E above).

The post-operative period was uneventful and an intra-uterine foleys catheter was inserted to keep the cavity patent. It was removed after 10 days. The patient was given conjugated estrogen for 21 days and medroxyprogesterone for last 10 days for 6 months.

DISCUSSION

Uterine bicornis was an asymptomatic incidental finding in course of radiographic studies of the uterus of this patient. Complimentary hysteroscopy review confirmed the earlier noted findings. There was also no tubal patency on HSG, which maybe as a result of the huge fibroids on the horns of the uterus. This is the usual pattern of arriving at a diagnosis for most cases of Bicornuate uterus and other congenital anomalies of the uterus.⁹⁻¹²

There are very few cases of fibroids co-existing with Mullerian anomalies reported in literature and thus the diagnosis is not often made because of the low incidence.¹³

Metroplastic surgery was described by Strassman in 1952 for class III, IV and V anomalies, and it was subsequently modified and simplified by Jones in 1953 (wedge excision of the septum) and Tompkins in 1962 (incision of the septum).^{4,14}

Open conventional metroplasty and laproscopy for the treatment of Bicornuate uterus are both safe and viable options.⁸ The patient had an abdominal modified strassman's metroplasty that involved excision of the septum. This procedure has been widely practiced in the few symptomatic cases of Bicornuate ueterus.^{4,8} Intra-operatively, in the hands of a skilled surgeon adequate care may be taken to ensure that the myometrial edges are not sutured under tension, as it is prone to hematoma formation.

In this patient laparoscopy was indicated as an option, but considering the multiple uterine fibroids, its size and unavailability of the facilities and experience, an open abdominal procedure was considered.

Post-operative hysterosalpingography studies confirmed a single cavity and patent tubes. Pregnancy has been widely reported following metroplasty, although there is increased risk of placenta previa, morbidly adherent placenta and uterine rupture.¹⁵⁻¹⁸

Considering the age of the patient, even though nulliparous, pregnancy outcome as recorded in literature holds a good prognosis for the patient.¹⁹

CONCLUSION

The correction of uterine anomalies is recommended in patients who show symptoms. Surgical metroplasty has been shown to be an effective method of treatment of the symptomatic patients and also offers improvement in fertility and pregnancy outcome.

The use of laparoscopic approach to myomectomy and metroplasty is gaining grounds worldwide and Africa need to rise up to the occasion in order to offer patients the benefits of these advancements in clinical practice.



REFERENCES

1. Saravelos SH, Cocksedge KA, Li TC. Prevalence and diagnosis of congenital uterine anomalies in women with reproductive failure: A critical appraisal. *Hum Reprod Update*. 2008;14(5):415–29.
2. Subburaj L, Rajkumar K. Laparoscopic metroplasty for bicornuate uterus – A case report. *Indian J Obstet Gynecol Res*. 2021;8(2):279–81.
3. Chandler TM, Machan LS, Cooperberg PL, Harris AC, Chang SD. Müllerian duct anomalies: From diagnosis to intervention. *Br J Radiol*. 2009;82(984):1034–42.
4. Passos I de MP e., Britto RL. Diagnosis and treatment of müllerian malformations. *Taiwan J Obstet Gynecol*. 2020;59(2):183–8.
5. Ribeiro SC, Tormena RA, Peterson TV, Gonzáles M de O, Serrano PG, de Almeida JAM, et al. Müllerian duct anomalies: Review of current management. *Sao Paulo Med J*. 2009;127(2):92–6.
6. Yadav A, Prateek S, Chawla L, Sharma S, Choudhary D. Bicornuate Uterus with Unilateral Fibroid - Surgical Procedure or LNG-IUS – A Conservative Approach in a Patient Who Opted LNG as Contraception. *J Evol Med Dent Sci*. 2020;9(39):2924–6.
7. Nwankwo NC, Maduforo CO. Mullerian duct anomaly in a Nigerian woman with recurrent pregnancy loss. *Niger J Clin Pract*. 2011;14(1):109–11.
8. Attaran M. Surgical Techniques for Management of Anomalies of the Müllerian Ducts. In: *Clinical Reproductive Medicine and Surgery*. 2017. p. 461–85.
9. Hg I, Sb M, Ar H, Ua U. Bicornuate Uterus Presenting with Recurrent Pregnancy Loss ; the Role of Hysterosalpingography in the Diagnosis : A Report of Two Cases. 2020;10(December):270–3.
10. Ludwin A, Ludwin I, Banas T, Knafel A, Miedzyblocki M. Diagnostic accuracy of sonohysterography , hysterosalpingography and diagnostic hysteroscopy in diagnosis of arcuate , septate and bicornuate uterus. 2011;37(3):178–86.
11. Article O. Comparative Prospective Study of Hysterosalpingography and Hysteroscopy in Infertile Women. 2017;
12. Tsonis O. Sonographic and Hysteroscopic Assessment of Uterine Congenital *Journal of*

- Medical - Clinical Research & Reviews
Sonographic and Hysteroscopic Assessment of Uterine Congenital Malformations: A Retrospective Study. 2021;(April):1–5.
13. Bafna B, Bafna A, Bafna A. Uterus bicornis unicollis with multiple leiomyomas. *Int J Reprod Contraception, Obstet Gynecol.* 2016;5(11):4088–90.
 14. Matsaseng T, Kruger TF. Laparoscopic Strassman's metroplasty for bicornuate uterus -is it relevant? *S Afr J Obstet Gynaecol.* 2012;18(3):85–7.
 15. Zhang C, Wang X, Jiang H, Hou L, Zou L. Placenta percreta after Strassman metroplasty of complete bicornuate uterus: a case report. *BMC Pregnancy Childbirth.* 2021;21(1):1–5.
 16. Zhou QY, Saravelos S, Huang XW, Ma N, Li T, Xia EL. Laparoscopic metroplasty for bicornuate uterus: surgical techniques and outcomes. *Chin Med J (Engl).* 2020;134(9):1107–9.
 17. Ono S, Yonezawa M, Watanabe K, Abe T, Mine K, Kuwabara Y, et al. Retrospective cohort study of the risk factors for secondary infertility following hysteroscopic metroplasty of the uterine septum in women with recurrent pregnancy loss. *Reprod Med Biol.* 2018;17(1):77–81.
 18. Gulavi E, Kyende Mutiso S, Mariara Muriuki C, Mukaindo Mwaniki A. Successful Pregnancy Outcome after Open Strassman Metroplasty for Bicornuate Uterus. *Case Rep Obstet Gynecol.* 2018;2018:1–4.
 19. Wang X, Hou H, Yu Q. Fertility and pregnancy outcomes following hysteroscopic metroplasty of different sized uterine septa: A retrospective cohort study protocol. *Med (United States).* 2019;98(30).

INDEX

Picture Gallery




(a) The Uterus with its two horns and fibroids in situ (b) After Reconstruction



View at hysteroscopy






Pre and post surgical HSG



Steps in surgical reconstruction



Final reconstructed uterus





Cite this Article: [Author] (2022). Bicornuate Uterus With Multiple Uterine Fibroids, Metroplasty and Myomectomy; Case Profile and Literature Review. *Greener Journal of Medical Sciences*, 12(1): 131-142.